

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
31 July 2008 (31.07.2008)

PCT

(10) International Publication Number
WO 2008/090198 A1

(51) International Patent Classification:

A61K 31/21 (2006.01) A61K 31/445 (2006.01)
 A61K 31/216 (2006.01) A61K 31/4468 (2006.01)
 A61K 31/4045 (2006.01) A61K 31/496 (2006.01)
 A61K 31/4155 (2006.01) A61P 9/10 (2006.01)

(21) International Application Number:

PCT/EP2008/050814

(22) International Filing Date: 24 January 2008 (24.01.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

07101162.1 25 January 2007 (25.01.2007) EP

(71) Applicant (for all designated States except US):
JANSSEN PHARMACEUTICA NV [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KING, Peter John** [GB/BE]; c/o Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE). **HRUPKA, Brian Joel** [US/BE]; c/o Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE). **BORGHYS, Herman Karel** [BE/BE]; c/o Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE). **BERWAER, Monique Jenny**Marie [BE/BE]; Route de Bergister, Grandménil 1, B-6960 Manhay (BE). **ROEVENS, Peter Walter Maria** [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(54) Title: USE OF MTP INHIBITORS FOR INCREASING LEVELS OF SATIETY HORMONES

(57) Abstract: The present invention relates to the use of inhibitors of microsomal triglyceride transfer protein (MTP) for increasing plasma levels of the satiety hormones such as GLP-1, PYY and CCK.

 WO 2008/090198 A1

USE OF MTP INHIBITORS FOR INCREASING LEVELS OF SATIETY HORMONES

5 **[0001]** The present invention relates to the use of inhibitors of microsomal triglyceride transfer protein (MTP) for increasing plasma levels of the satiety hormones such as GLP-1, PYY and CCK.

10 **[0002]** Microsomal triglyceride transfer protein (hereinafter referred to as MTP) is known to catalyze the transport of triglyceride, cholesteryl ester and phospholipids such as phosphatidylcholine. This indicates that MTP is required for the synthesis of Apo B-containing lipoproteins such as chylomicrons and VLDL, the precursor to LDL. It therefore follows that an MTP inhibitor would inhibit the synthesis of VLDL and chylomicrons, thereby lowering levels of VLDL, LDL, cholesterol and triglyceride in humans. Compounds capable of inhibiting MTP are believed to be useful in the
15 treatment of disorders such as obesity, hyperlipidemia, hypercholesterolemia, hypertriglyceridemia, class II diabetes, atherosclerosis and for the reduction of postprandial serum triglyceride plasma levels.

20 **[0003]** Satiety hormones are hormones released from the gastrointestinal tract in response to changes in the nutritional state. These hormones influence central mechanisms involved in the regulation of energy balance, through a range of bloodborne and neural pathways.

25 **[0004]** Glucagon-like peptide 1 (GLP-1) is an intestinal hormone which generally stimulates insulin secretion during hyperglycemia, suppresses glucagon secretion, stimulates (pro) insulin biosynthesis and decelerates gastric emptying and acid secretion. GLP-1 is secreted from L cells in the small and large bowel following the ingestion of fat and proteins. GLP-1 has been implicated as a possible therapeutic agent for the management of type 2 non-insulin-dependent diabetes mellitus as well
30 as related metabolic disorders, such as obesity.

35 **[0005]** Pancreatic polypeptide ("PP") was discovered as a contaminant of insulin extracts and was named by its organ of origin rather than functional importance. A related peptide was subsequently discovered in extracts of intestine and named Peptide YY ("PYY") because of the N- and C-terminated tyrosines (Tatemoto, Proc. Natl. Acad. Sci. USA, 79 : 2514 –2518 (1982)). PYY is secreted from the endocrine L cells of the small and large bowel, with high concentration at the terminal ileum, colon and maximum concentration in the rectum. Plasma PYY levels are suppressed in the

- 2 -

fasted state and increase within 30 minutes of nutrients reaching the gut. PYY release is stimulated by nutrient intake in proportion to energy content. It is particularly stimulated by fat intake, compared to carbohydrate and protein meals with a similar calorie content. Recent studies suggest that PYY can induce appetite reduction.

5

[0006] Cholecystikinin is structurally related to gastrin and exists in several molecular forms with differing numbers of amino acids – examples include CCK-8, CCK-33, CCK-39 and CCK-54. CCK is an endogenous gut hormone found mainly within the duodenum and jejunum and is released following the consumption of food.

10 Release of CCK has been shown to be a satiety signal in humans. When food is consumed, CCK releasing protein (CCKRP) is released in the small intestine. CCKRP stimulates CCK release from the intestinal cells. It has been shown that CCK release results in appetite reduction so that the person will stop eating.

15 **[0007]** The ability of CCK to reduce appetite appears to make it a useful agent in the treatment of obesity. An increase in the level of the satiety hormone CCK would result in less food consumed and reduction of hunger cravings between meals. These effects would enable an overweight individual to better comply with a diet that has a reduced caloric intake.

20

[0008] An increase in the level of the satiety hormone CCK extends the feeling of satiety, resulting in a decrease of food intake which over time results in a decrease in body weight while providing better regulation of glucose and insulin levels following consumption of a meal. The release of CCK also causes a delay in stomach emptying which blunts the post-prandial rise in glucose and insulin. Most persons with Type II

25 diabetes are obese and have an inability to respond normally to insulin. An increase in CCK levels may permit Type II diabetics to be satiated with a lower caloric intake and may offer a better degree of glycemic control.

30 **[0009]** Bulimia is an eating disorder characterised by an inability to become satiated by food. As a result bulimics tend to binge on food and regurgitate it to prevent weight gain. Studies have shown that bulimics have a defect in their normal satiety mechanism. Hence an increase of the satiety hormones would permit bulimics to feel satiated.

35

[0010] Unexpectedly it has now been observed that when inhibitors of microsomal triglyceride transfer protein (MTP) are administered to a mammalian subject, the plasma levels of the satiety hormones such as GLP-1, PYY and CCK are increased.

- 3 -

[0011] The present invention provides the use of a MTP inhibiting compound for the manufacture of a medicament for increasing the levels of satiety hormones, such as the GLP-1, PYY and CCK hormones. Also provided is the use of a pharmaceutical composition comprising a MTP inhibiting compound for the manufacture of a
5 medicament for increasing the levels of satiety hormones, such as the GLP-1, PYY and CCK hormones.

[0012] Further, the present invention provides a method for increasing the levels of satiety hormones, in particular GLP-1, PYY and CCK, in a mammalian subject, which
10 method comprises administering to a mammal a therapeutically effective amount of an MTP inhibiting compound or a pharmaceutical composition comprising a MTP inhibiting compound.

[0013] The use of MTP inhibiting compound for increasing the levels of satiety
15 hormones, in particular the GLP-1, PYY and CCK hormones, also has a lowering effect on the level of glucose in blood plasma and increases insulin sensitivity. Insulin resistance is the condition in which normal amounts of insulin are inadequate to produce a normal insulin response from fat, muscle and liver cells. Insulin resistance in fat cells results in hydrolysis of stored triglycerides, which elevates free fatty acids in
20 the blood plasma. Insulin resistance in muscle reduces glucose uptake whereas insulin resistance in liver reduces glucose storage, with both effects serving to elevate blood glucose. High plasma levels of insulin and glucose due to insulin resistance often leads to the metabolic syndrome and type 2 diabetes.

[0014] Studies in dogs with an induced dilated cardiomyopathy have shown that a
25 48 hour of GLP-1 infusion improved the left ventricular function, and reduced systemic vascular resistance compared with saline-treated control animals (Nikolaidis LA et al., Circulation 2004;110:955-961). Accordingly the present invention also relates to the use of MTP inhibiting compounds for increasing the levels of the satiety hormone
30 GLP-1 for the treatment of cardiomyopathy.

[0015] Studies in rats with pyridoxine induced peripheral sensory neuropathy suggest neuroprotection mediated by agonism at the GLP-1 receptor (Perry T. et al, Experimental Neurology 2007:203, 293 – 301). Accordingly the present invention also
35 relates to the use of MTP inhibiting compounds for increasing the levels of the satiety hormone GLP-1 for the treatment of peripheral neuropathies.

- 4 -

[0016] MTP inhibiting compounds have been disclosed in, e.g., Janssen Pharmaceutica : WO-96/13499, WO-02/20501, WO-02/42271, WO-02/081460, WO-2005/058824, and WO-2005/085226; Bristol-Myers-Squibb : EP-0,584,446, EP-0,643,057, WO-96/26205, WO-97/26240, WO-91/43255, WO-97/43257, 5 WO-98/27979, and WO-99/21564; GSK : WO-98/16526, WO-98/47877, WO-98/56790, WO-00/32582, WO-01/92241, WO-01/96327, and WO-03/048121; Japan Tobacco : WO-99/31085, WO-03/072532, and WO-2006/008962; Meiji Seika Kaisho : WO-98/54135; Novartis : WO-01/77077 and WO-2000/005201; Pfizer : WO-96/40640, and WO-98/23593.

10

[0017] Particular MTP inhibiting compounds are, e.g., dirlotapide or (S)-N-{2-[benzyl(methyl)amino]-2-oxo-1-phenylethyl}-1-methyl-5-[4'-(trifluoromethyl)[1,1'-biphenyl]-2-carboxamido]-1H-indole-2-carboxamide; BMS201038 or N-(2,2,2-trifluoroethyl)-9-[4-[4-[[4'-(trifluoromethyl)-1,1'-biphenyl-2-yl]carbonyl]amino)-1-15 piperidinyl]butyl]-9H-fluorene-9-carboxamide (EP-0,643,057); mitratapide or (-)-[2S-[2 α ,4 α (S*)]]-4-[4-[4-[4-[[2-(4-chlorophenyl)-2-[[4-methyl-4*H*-1,2,4-triazol-3-yl]thio]methyl]-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-2,4-dihydro-2-(1-methylpropyl)-3*H*-1,2,4-triazol-3-one (WO-96/13499); (+)-phenyl-(4-{4-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-phenyl}-piperidin-1-yl)-acetic acid methyl 20 ester (WO-02/20501); JTT-130 or diethyl ester[[[3-[(dimethylamino)carbonyl]-4-[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]acetyl]oxy]methyl]phenyl propanedioic acid (WO-2006/008962); SLx 4090 from Surface Logix; NA-2003 from Meiji Seika Kaisha; [(2*R*)-2,3-dihydro-5-[[[6-methyl-4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]-1H-inden-2-yl]-carbamic acid methyl ester (WO-2000/005201); 25 T-0126 or N-[2-[2-(1*H*-Pyrazol-1-yl)acetyl]-2,3-dihydro-1*H*-isoindol-5-yl]-2-[5-(trifluoromethyl)pyridin-2-yl]benzamide from Tanabe Seiyaku (WO-2002/014276).

30

[0018] As used herein, "mammal" or "mammalian subject" refers to human and non-human patients.

[0019] As used herein, a "therapeutically effective amount" of a MTP inhibiting compound, is the quantity of a compound which, when administered to a mammalian subject, results in a sufficiently high level of that MTP inhibiting compound in the mammalian to cause a discernible increase of the blood plasma levels of the satiety

- 5 -

hormones GLP-1, PYY and CCK.

[0020] The pharmaceutical compositions comprising a MTP inhibiting compound can be administered to a subject either orally, parenterally (for example intravenously, 5 intramuscularly or subcutaneously), percutaneously, or rectally.

[0021] Solid dosage forms for oral administration include capsules, dragees, tablets, powders and granules. These solid dosage forms are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as 10 used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined amount of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, 15 teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

[0022] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, suspo-emulsions, syrups and elixirs. Pharmaceutical compositions for parenteral injection may comprise physiologically 20 acceptable sterile aqueous or nonaqueous solutions, dispersions, suspension, or emulsions, or may comprise sterile powders for reconstitution into sterile injectable solutions or dispersions.

Description of the drawings

25 **[0023]** Figure 1 is a graph displaying plasma CCK (pMol/l) expressed as the median value per group just before the meal and 12 and 24 hours after the meal.

[0024] Figure 2 is a graph displaying the postprandial plasma PYY (pMol/l) levels after 0.02% (w/w) administration of the MTP inhibitor "compound A" mixed in diet 30 containing 17.5% (w/w) (35 kcal%) fat.

[0025] Figure 3 is a graph displaying the postprandial plasma GLP-1 (pg/l) levels after 0.02% (w/w) administration of the MTP inhibitor "compound A" mixed in diet containing 17.5% (w/w) (35 kcal%) fat.

Experimental part**Experiment 1 : Plasma CCK levels – single dose study in dog**

[0026] The effect of the MTP inhibitor (+)-phenyl-(4-{4-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-phenyl}-piperidin-1-yl)-acetic acid methyl ester (WO-02/20501)
5 (hereinafter referred to as “compound A”) on CCK plasma levels was studied in
3 groups of 8 dogs each (4 male and 4 female dogs per group). Two groups were
treated orally with two different doses of compound A and one group was treated orally
with the vehicle and served as a placebo group. The vehicle solution contained the
same ingredients as the test formulations with omission of the test substance
10 compound A.

[0027] The treatment groups were :

- group (1) treated orally with vehicle
- group (2) treated orally with 0.15 mg compound A per kg body weight
- 15 - group (3) treated orally with 0.63 mg compound A per kg body weight

[0028] Dosing with either vehicle (group 1) or compound A (groups 2 and 3) was
done together with a liquid meal at 7.00 hour for the first two males and females of
each group and at 18.30 h for the last two males and females of each group. CCK
20 plasma levels were determined before dosing, 0 hour, and at 12 and 24 hours post
feeding.

[0029] As seen in Figure 1, plasma CCK (pMol/l) expressed as the median value per
group just before the meal and 12 and 24 hours after the meal showed a dose related
25 increase of plasma CCK levels after administration of the MTP inhibiting compound A.

Experiment 2 : Plasma PYY levels – study in rats

[0030] Male Sprague-Dawley rats (Iffa-Credo) are housed in individually ventilated
cages under controlled temperature (20-24°C), humidity (45-65%) and light (12-12h
30 light/dark cycle; Lights on - 5 AM – 5PM). Rats were adapted to a semipurified casein,
cornstarch and sucrose based diet (AIN-93) containing 17.5% w/w corn oil as the fat
source for 10 days. The 17.5% diet is calculated to contain 35% of energy as fat.

[0031] At dark onset on day 11, half the rats were switched to the same diet
35 containing 0.02% w/w of “compound A”, while the remaining rats received the
control/adaptation diet. At 0, 1, 2, 4, 6, 12, 14, 16, 20 and 24 hr after diet presentation,

- 7 -

a group of 6 rats/treatment were killed by decapitation and 4 ml of trunk blood was collected in pre-cooled (4°C) K3E plasma tubes containing protease inhibitor cocktail. Blood was centrifuged (1500 x g for 15 minutes at 4°C) within 10-15 minutes of sample collection taking blood sample and stored at -70°C until assayed.

5

Experiment 3 : Plasma GLP-1 levels – study in rats

[0032] Male Sprague-Dawley rats (Iffa-Credo) are housed in individually ventilated cages under controlled temperature (20-24°C), humidity (45-65%) and light (12-12h light/dark cycle; Lights on - 5 AM – 5PM). Rats were adapted to a semipurified casein, cornstarch and sucrose based diet (AIN-93) containing 17.5% w/w corn oil as the fat source for 10 days. The 17.5% diet is calculated to contain 35% of energy as fat.

[0033] At dark onset on day 11, half the rats were switched to the same diet containing 0.02% w/w of “compound A”, while the remaining rats received the control/adaptation diet. At 0, 1, 2, 4, 6, 12, 14, 16, 20 and 24 hr after diet presentation, a group of 6 rats/treatment were killed by decapitation and 4 ml of trunk blood was collected in pre-cooled (4°C) K3E plasma tubes containing protease inhibitor cocktail. Blood was centrifuged (1500 x g for 15 minutes at 4°C) within 10-15 minutes of sample collection taking blood sample and stored at -70°C until assayed.

20

Claims

1. Use of a MTP inhibiting compound for the manufacture of a medicament for the treatment of a disease mediated by increasing the levels of satiety hormones.
- 5 2. Use according to claim 1 wherein the satiety hormones are GLP-1, PYY and CCK.
3. Use according to claim 2 wherein the satiety hormone is GLP-1.
4. Use according to claim 2 wherein the satiety hormone is PYY.
- 10 5. Use according to claim 2 wherein the satiety hormone is CCK.
6. Use of a MTP inhibiting compound for the manufacture of a medicament for increasing the levels of the satiety hormones GLP-1, PYY and CCK and concomitant lowering of glucose levels.
- 15 7. Use of a MTP inhibiting compound for the manufacture of a medicament for increasing the levels of the satiety hormones GLP-1, PYY and CCK and concomitant lowering of insulin sensitivity.
- 20 8. The use as claimed in claim 2 wherein the disease is cardiomyopathy.
9. The use as claimed in claim 2 wherein the disease is peripheral neuropathies.
- 25 10. The use according to any of claims 1 to 9 wherein the MTP inhibiting compound is selected from dirlotapide , N-(2,2,2-trifluoroethyl)-9-[4-[4-[(4'-trifluoromethyl)-1,1'-biphenyl-2-yl]carbonyl]amino)-1-piperidinyl]butyl]-9H-fluorene-9-carboxamide; (-)-[2S-[2 α ,4 α (S*)]]-4-[4-[4-[4-[[2-(4-chlorophenyl)-2-[[4-methyl-4*H*-1,2,4-triazol-3-yl)thio]methyl]-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-2,4-
- 30 dihydro-2-(1-methylpropyl)-3*H*-1,2,4-triazol-3-one; (+)-phenyl-(4-{4-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-phenyl}-piperidin-1-yl)-acetic acid methyl ester; diethyl ester[[[3-[(dimethylamino)carbonyl]-4-[[4'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl]carbonyl]amino]phenyl]acetyl]oxy]methyl]phenyl propanedioic acid; [(2*R*)-2,3-dihydro-5-[[[6-methyl-4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]-carbonyl]amino]-1*H*-inden-2-yl]-carbamic acid methyl ester (WO-2000/005201); or
- 35 N-[2-[2-(1*H*-pyrazol-1-yl)acetyl]-2,3-dihydro-1*H*-isoindol-5-yl]-2-[5-(trifluoromethyl)pyridin-2-yl]benzamide.
- 40 11. The use according to claim 10 wherein the MTP inhibiting compound is (-)-[2S-[2 α ,4 α (S*)]]-4-[4-[4-[4-[[2-(4-chlorophenyl)-2-[[4-methyl-4*H*-1,2,4-triazol-3-

- 9 -

yl)thio]methyl]-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-2,4-dihydro-2-(1-methylpropyl)-3*H*-1,2,4-triazol-3-one.

12. The use according to claim 10 wherein the MTP inhibiting compound is
- 5 (+)-phenyl-(4-{4-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-phenyl}-piperidin-1-yl)-acetic acid methyl ester.

5 **Figure 1** : Plasma CCK (pMol/l) expressed as the median value per group just before the meal and 12 and 24 hours after the meal. Dosing of the MTP inhibitor “compound A” was done together with the meal which consisted of a liquid meal given orally by gavage.





Figure 2 : Postprandial Plasma PYY (pMol/l) levels after 0.02% (w/w) administration of the MTP inhibitor “compound A” mixed in diet containing 17.5% (w/w) fat

5





Figure 3 : Postprandial Plasma GLP-1 (pg/l) levels after 0.02% (w/w) administration of the MTP inhibitor “compound A” mixed in diet containing 17.5% (w/w) fat

5



INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2008/050814

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K31/21 A61K31/216 A61K31/4045 A61K31/4155 A61K31/445
A61K31/4468 A61K31/496 A61P9/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 1 099 438 A (PFIZER PROD INC [US]) 16 May 2001 (2001-05-16) abstract page 1, paragraph 4 - paragraph 6 page 3, paragraph 10 page 12, paragraph 105	1-7, 10
X	WO 2005/046644 A (PFIZER PROD INC [US]; FRIESEN DWAYNE THOMAS [US]; SHANKER RAVI MYSORE) 26 May 2005 (2005-05-26) page 1, line 5 - line 8 page 2, line 15 - line 21 page 4, line 17	1-7, 10
	----- -/--	

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

18 June 2008

Date of mailing of the international search report

25/06/2008

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Damiani, Federica

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2008/050814

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 712 279 A (BILLER SCOTT A [US] ET AL) 27 January 1998 (1998-01-27) column 1, line 15 - line 18 column 23 column 40, line 25 - line 34 -----	1-7, 10
X	WO 00/05201 A (NOVARTIS AG [CH]; NOVARTIS ERFIND VERWALT GMBH [AT]; FINK CYNTHIA ANNE) 3 February 2000 (2000-02-03) cited in the application page 20, paragraph 3 page 84; example 14 -----	1-8, 10
X	JP 2003 231635 A (TANABE SEIYAKU CO) 19 August 2003 (2003-08-19) abstract -----	1-7, 10
X	WO 2006/008962 A (JAPAN TOBACCO INC [JP]; FURUKAWA NOBORU [JP]; MERA YASUKO [JP]; KAWAI) 26 January 2006 (2006-01-26) cited in the application abstract -----	1-7, 10
X	WO 2005/070390 A (JANSSEN PHARMACEUTICA NV [BE]; FRANCOIS MARC KAREL JOZEF [BE]; EMBRECH) 4 August 2005 (2005-08-04) page 1, paragraphs 1,2 page 7, paragraph 25 -----	1-8, 10, 11
X	WO 02/20501 A (JANSSEN PHARMACEUTICA NV [BE]; MEERPOEL LIEVEN [BE]; ROEVENS PETER WAL) 14 March 2002 (2002-03-14) cited in the application example B.10; tables F-1 page 25, line 1 - line 8 page 25, line 24 - line 30 -----	1-7, 10, 12
X	WO 00/53171 A (MOLTENI L E C DEI FRATELLI ALI [IT]; MANNUCCI EDOARDO [IT]; ROTELLA CA) 14 September 2000 (2000-09-14) page 1, line 5 - line 8 page 8 -----	1-7
X	WO 2004/022074 A1 (NOVARTIS NUTRITION AG [CH]; HUGHES THOMAS EDWARD [US]; WESTERTERP-PLAN) 18 March 2004 (2004-03-18) abstract page 1, line 1 - line 4 page 3, line 5 - line 9 page 2, line 26 -----	1-7, 9

-/--

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2008/050814

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2006/204567 A1 (HU JIANG [US] ET AL) 14 September 2006 (2006-09-14) abstract page 1, paragraph 5 -----	1-7
X	EP 1 685 834 A (LODERS CROKLAAN BV [NL]) 2 August 2006 (2006-08-02) page 3, paragraph 16 page 9, paragraph 66 -----	1-7
X	WO 2004/047755 A (TULARIK INC [US]; JAPAN TOBACCO INC [JP]; FOX BRIAN M [US]; FURUKAWA N) 10 June 2004 (2004-06-10) claims 27,30 -----	1-8
X	WO 03/092694 A (PFIZER PROD INC [US]; TRACEY WAYNE ROSS [US]; TREADWAY JUDITH LEE [US]) 13 November 2003 (2003-11-13) abstract page 25, line 9 - line 11 claims 3,8 -----	1-9
P,X	US 2007/093527 A1 (WISLER GERALD L [US]) 26 April 2007 (2007-04-26) abstract page 3, paragraph 27 -----	1-7,9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2008/050814

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1099438	A	16-05-2001	AU 777542 B2	21-10-2004
			AU 7151900 A	17-05-2001
			CA 2325282 A1	10-05-2001
			HU 0004454 A2	28-10-2001
			JP 2001181209 A	03-07-2001
			KR 20010051509 A	25-06-2001
			NZ 508061 A	26-04-2002
			ZA 200006417 A	08-05-2002
WO 2005046644	A	26-05-2005	AR 048206 A1	12-04-2006
			AU 2004289110 A1	26-05-2005
			BR PI0416596 A	30-01-2007
			CA 2545443 A1	26-05-2005
			CN 1878538 A	13-12-2006
			EP 1696887 A1	06-09-2006
			JP 2007511500 T	10-05-2007
			KR 20060096076 A	05-09-2006
			MX PA06005489 A	11-08-2006
US 5712279	A	27-01-1998	AR 001795 A1	10-12-1997
			AT 283851 T	15-12-2004
			AU 699865 B2	17-12-1998
			AU 4763196 A	11-09-1996
			BG 101717 A	27-02-1998
			CA 2213466 A1	29-08-1996
			CN 1176640 A	18-03-1998
			CZ 9702617 A3	14-01-1998
			DE 69633983 D1	05-01-2005
			DE 69633983 T2	22-12-2005
			EE 9700182 A	16-02-1998
			EP 0886637 A1	30-12-1998
			ES 2233961 T3	16-06-2005
			FI 973416 A	20-08-1997
			HU 9801278 A2	28-06-1999
			IL 116917 A	31-08-2000
			JP 4036244 B2	23-01-2008
			JP 11500442 T	12-01-1999
			LV 11951 A	20-01-1998
			NO 973821 A	20-08-1997
			NZ 302055 A	28-02-2000
			PL 322003 A1	05-01-1998
			SK 113597 A3	09-09-1998
			TW 486469 B	11-05-2002
			WO 9626205 A1	29-08-1996
			UY 24165 A1	31-07-2001
WO 0005201	A	03-02-2000	AR 029447 A1	02-07-2003
			AU 5161399 A	14-02-2000
			CA 2338198 A1	03-02-2000
			EP 1097129 A1	09-05-2001
			JP 2002521360 T	16-07-2002
JP 2003231635	A	19-08-2003	NONE	
WO 2006008962	A	26-01-2006	AU 2005264301 A1	26-01-2006
			BR PI0513368 A	06-05-2008
			CA 2574006 A1	26-01-2006
			EP 1769793 A1	04-04-2007

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2008/050814

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2006008962 A		KR 20070036126 A	02-04-2007
WO 2005070390 A	04-08-2005	AU 2005205933 A1 CA 2552988 A1 JP 2007518774 T	04-08-2005 04-08-2005 12-07-2007
WO 0220501 A	14-03-2002	AU 1046802 A BG 107581 A BR 0114045 A CA 2421228 A1 CN 1449382 A CZ 20030892 A3 EA 5855 B1 EE 200300080 A HK 1057895 A1 HR 20030156 A2 HU 0302230 A2 IS 6682 A JP 2004508361 T MX PA03001890 A NO 20031001 A NZ 524525 A PL 366085 A1 SK 3952003 A3 UA 77946 C2 US 2004014971 A1 ZA 200301755 A	22-03-2002 28-11-2003 01-07-2003 14-03-2002 15-10-2003 18-02-2004 30-06-2005 15-02-2005 26-10-2007 30-04-2003 28-10-2003 15-01-2003 18-03-2004 24-06-2003 04-03-2003 26-03-2004 24-01-2005 03-02-2004 15-02-2007 22-01-2004 22-06-2004
WO 0053171 A	14-09-2000	AU 3960400 A	28-09-2000
WO 2004022074 A1	18-03-2004	AU 2003260495 A1 EP 1536805 A1 JP 2006508057 T	29-03-2004 08-06-2005 09-03-2006
US 2006204567 A1	14-09-2006	NONE	
EP 1685834 A	02-08-2006	NONE	
WO 2004047755 A	10-06-2004	AU 2003293006 A1 BR 0315688 A CA 2514473 A1 CN 1753897 A EP 1562956 A2 JP 3988830 B2 JP 2006509764 T KR 20050090986 A MX PA05005425 A NZ 539952 A ZA 200503823 A	18-06-2004 06-09-2005 10-06-2004 29-03-2006 17-08-2005 10-10-2007 23-03-2006 14-09-2005 23-11-2005 30-05-2008 22-02-2006
WO 03092694 A	13-11-2003	AU 2003219421 A1 BR 0309707 A CA 2483927 A1 EP 1499317 A1 MX PA04008646 A	17-11-2003 09-02-2005 13-11-2003 26-01-2005 06-12-2004
US 2007093527 A1	26-04-2007	NONE	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2008/050814

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
<hr/>			